

A01-O10

DETECTION OF TRENDS IN CO₂ VERTICAL PROFILES DUE TO BIOSPHERIC AND ANTHROPOGENIC FLUX CHANGE

Prabir K. Patra (*Department of Environmental Geochemical Cycle Research, JAMSTEC, Japan / Center for Atmospheric and Oceanic Studies, Tohoku University, Japan*)

Toshinobu Machida (*National Institute for Environmental Studies (NIES), Japan*)

Shuji Aoki (*Center for Atmospheric and Oceanic Studies, Tohoku University, Japan*)

Tazu Saeki, (*Department of Environmental Geochemical Cycle Research, JAMSTEC, Japan*)

Takakiyo Nakazawa (*Center for Atmospheric and Oceanic Studies, Tohoku University, Japan*)

prabir@jamstec.go.jp

Large increase in emissions of CO₂ has been estimated from fossil fuel consumption statistics since the 1990 as per the CDIAC, IEA or EDGAR. Independent studies also propose that the terrestrial biospheric uptake must have been increased in order to balance the sources, sinks and atmospheric burden. However, the relative sources and sinks strengths still have large uncertainties due to lack of sufficient observations and uncertainties in modeling tools. Since the early 1990s, vertical profiles of atmospheric CO₂ are measured by the both NIES (in Siberia) and Tohoku University (over Japan) observational programs. We use these vertical profile observations and an AGCM-based chemistry transport model (JAMSTEC's ACTM) to decipher the signals from fossil fuel consumption, using measurements over Japan that samples the outflow of the continental emissions, and terrestrial biosphere of the arctic, which has supposedly ventilating greater amount of carbon due to increased surface temperature and CO₂ fertilization.

The ACTM simulation results, using the fluxes from a 84-region time-dependent inverse model and increasing fossil fuel emissions, suggest that the CO₂ vertical gradients should have become greater over Japan during the winter as the fossil fuel CO₂ signal from the continental Asia outflows isentropically near the surface. However, during the summer the continental outflow occur through the upper troposphere, resulting in weaker vertical gradient or even inverted vertical profile over Japan. The ACTM results over Surgut and Nobosibirisk (NIES sites in Siberia) suggest that the terrestrial biospheric uptake increase consists of detectable change in the gradients in CO₂ vertical profiles. These results, in comparison with measured vertical profiles using research aircrafts, will be presented.